Note On Elliptic Groups of Prime Orders

N. A. Carella

***Abstract*:** Let E be an elliptic curve of rank $\mathrm{rk}(E) \geq 0$, and let $E(\mathbb{F}_p)$ be the elliptic group of order $\#E(\mathbb{F}_p) = n$. The number of primes $p \leq x$ such that n is prime is expected to be $\pi(x, E) = \delta(E)x/\log^2 x + o(x/\log^2 x)$, where $\delta(E) \geq 0$ is the density constant. This note proves a lower bound $\pi(x, E) \gg x/\log^3 x$.

Mathematics Subjects Classification: Primary 11G07; Secondary 11N36, 11G15.
Keywords: Elliptic Prime; Group of Prime Order, Primitive Point; Koblitz Conjecture.

arXiv:1702.06814v2 [math.GM] 11 Apr 2017

1 Introduction

The applications of elliptic curves in cryptography demands elliptic groups of certain orders $n = \#E(\mathbb{F}_p)$, and certain factorizations of the integers n. The extreme cases have groups of $\mathbb{F}_p$-rational points $E(\mathbb{F}_p)$ of prime orders, and small multiples of large primes.

Conjecture 1.1 (Koblitz)**.** *Let $E : f(X, Y) = 0$ be an elliptic curve of of discriminant $\Delta \neq 0$ defined over the integers $\mathbb{Z}$ which is not $\mathbb{Q}$-isogenous to a curve with nontrivial $\mathbb{Q}$-torsion and does not have CM. Then,*

$$\begin{aligned} \pi(x, E) &= \#\{p \leq x : p \not| \Delta \text{ and } \#E(\mathbb{F}_p) = \text{prime}\} \\ &= \frac{x}{\log^2 x} \prod_{p \geq 2} \left(1 - \frac{p^2 - p - 1}{(p-1)^3(p+1)}\right) + O\left(\frac{x}{\log^3 x}\right), \end{aligned} \tag{1}$$

for large $x \geq 1$.

The product expression appearing in the above formula is basically the average density of prime orders, some additional details are given in Section 7. A result for groups of prime orders generated by primitive points is proved here. Let

$$\pi(x, E) = \#\{p \leq x : p \not| \Delta \text{ and } d_E^{-1} \cdot \#E(\mathbb{F}_p) = \text{prime}\}. \tag{2}$$

The parameter $d_E \geq 1$ is a small integer defined in Section 6.

Theorem 1.1. *Let $E : f(X, Y) = 0$ be an elliptic curve over the rational numbers $\mathbb{Q}$ of rank $\mathrm{rk}(E(\mathbb{Q}) > 0$. Then, as $x \to \infty$,*

$$\pi(x, E) \geq \delta(d_E, E) \frac{x}{\log^3 x} \left(1 + O\left(\frac{x}{\log x}\right)\right), \tag{3}$$

where $\delta(d_E, E)$ is the density constant.

The proof of this result is split into several parts. The next sections are intermediate results. The proof of Theorem 1.1 is assembled in the penultimate section, and the last section has examples of elliptic curves with infinitely many elliptic groups $E(\mathbb{F}_p)$ of prime orders n.

2 Representation of the Characteristic Function

2.1 Primitive Points Tests

For a prime $p \geq 2$, the group of points on an elliptic curve $E : y^2 = f(x)$ is denoted by $E(\mathbb{F}_p)$. Several definitions and elementary properties of elliptic curves and the n-division polynomial $\psi_n(x, y)$ are sketched in Chapter 14.

Definition 2.1. The order $\min\{k \in \mathbb{N} : kP = \mathcal{O}\}$ of an elliptic point is denoted by $\mathrm{ord}_E(P)$. A point is a *primitive point* if and only if $\mathrm{ord}_E(P) = n$.

Lemma 2.1. *If $E(\mathbb{F}_p)$ is a cyclic group, then it contains a primitive point $P \in E(\mathbb{F}_p)$.*

Proof. By hypothesis, $E(\mathbb{F}_p) \cong \mathbb{Z}/n\mathbb{Z}$, and the additive group $\mathbb{Z}/n\mathbb{Z}$ contains $\varphi(n) \geq 1$ generators (primitive roots). ■

More generally, there is map into a cyclic group

$$E(\mathbb{Q})/E(\mathbb{Q})_{tors} \longrightarrow \mathbb{Z}/m\mathbb{Z}, \tag{4}$$

for some $m = \#E(\mathbb{F}_p)/d$, with $d \geq 1$; and the same result stated in the Lemma holds in the smaller cyclic group $\mathbb{Z}/m\mathbb{Z} \subset \mathbb{Z}/n\mathbb{Z}$.

Lemma 2.2. *Let $\#E(\mathbb{F}_p) = n$ and let $P \in E(\mathbb{F}_p)$. Then, P is a primitive point if and only if $(n/q)P \neq \mathcal{O}$ for all prime divisors $q \,|\, n$.*

Basically, this is the classical Lucas-Lehmer primitive root test applied to the group of points $E(\mathbb{F}_p)$. Another primitive point test intrinsic to elliptic curves is the n-division polynomial test.

Lemma 2.3. (n-Division primitive point test) *Let $\#E(\mathbb{F}_p) = n$ and let $P \in E(\mathbb{F}_p)$. Then, P is a primitive point if and only if $\psi_{n/q}(P) \neq 0$ for all prime divisors $q \,|\, n$.*

The basic proof stems from the division polynomial relation

$$mP = \mathcal{O} \Longleftrightarrow \psi_m(P) = 0, \tag{5}$$

see [28, Proposition 1.25]. The elliptic primitive point test calculations in the penultimate lemma takes place in the set of integer pairs $\mathbb{Z} \times \mathbb{Z}$, while the calculations for the n-division polynomial primitive point test takes place over the set of integers $\mathbb{Z}$. The elementary properties of the n-division polynomials are discussed in [28], and the periodic property of the n-division polynomials appears in [26].

2.2 Additive Elliptic Character

The discrete logarithm function, with respect to the fixed primitive point T, maps the group of points into a cyclic group. The diagram below specifies the basic assignments.

$$\begin{array}{lcl} E(\mathbb{F}_p) & \longrightarrow & \mathbb{Z}/n\mathbb{Z}, \\ \mathcal{O} & \longrightarrow & \log_T(\mathcal{O}) = 0, \\ T & \longrightarrow & \log_T(T) = 1. \end{array} \tag{6}$$

In view of these information, an important character on the group of $E(\mathbb{F}_p)$-rational points can be specified.

Definition 2.2. A nontrivial additive elliptic character χ mod n on the group $E(\mathbb{F}_p)$ is defined by

$$\chi(\mathcal{O}) = e^{\frac{i2\pi}{n}\log_T(O)} = 1, \tag{7}$$

and

$$\chi(mT) = e^{\frac{i2\pi}{n}\log_T(mT)} = e^{i2\pi m/n}, \tag{8}$$

where $\log_T(mT) = m$ with $m \in \mathbb{Z}$.

2.3 Divisors Dependent Characteristic Function

A characteristic function for primitive points on elliptic curve is described in the [27, p. 5]; it was used there to derive a primitive point search algorithm.

Lemma 2.4. *Let E be a nonsingular elliptic curve, and let $E(\mathbb{F}_p)$ be its group of points of cardinality $n = \#E(\mathbb{F}_p)$. Let χ be the additive character of order d modulo d, and assume $P = (x_0, y_0)$ is a point on the curve. Then,*

$$\Psi_E(P) = \sum_{d\,|\,n} \frac{\mu(d)}{d} \sum_{0\le t<d} \chi(tP) = \begin{cases} 1 & \text{if } \mathrm{ord}_E(P) = n, \\ 0 & \text{if } \mathrm{ord}_E(P) \neq n. \end{cases} \tag{9}$$

2.4 Divisors Free Characteristic Function

A new *divisors-free* representation of the characteristic function of elliptic primitive points is developed here. This representation can overcomes some of the limitations of its equivalent in 2.4 in certain applications. The *divisors representation* of the characteristic function of elliptic primitive points, Lemma 2.4, detects the order $\mathrm{ord}_E(P)$ of the point $P \in E(\mathbb{F}_p)$ by means of the divisors of the order $n = \#E(\mathbb{F}_p)$. In contrast, the *divisors-free representation* of the characteristic function, Lemma 2.5, detects the order $\mathrm{ord}_E(P) \ge 1$ of a point $P \in E(\mathbb{F}_p)$ by means of the solutions of the equation $mT - P = \mathcal{O}$, where $P, T \in E(\mathbb{F}_p)$ are fixed points, $\mathcal{O}$ is the identity point, and m is a variable such that $0 \le m \le n-1$, and $\gcd(m, n) = 1$.

Lemma 2.5. *Let $p \geq 2$ be a prime, and let T be a primitive point in $E(\mathbb{F}_p)$. For a nonzero point $P \in E(\mathbb{F}_p)$ of order n the following hold: If $\chi \neq 1$ is a nonprincipal additive elliptic character of order* $\operatorname{ord}\chi = n$*, then*

$$\Psi_E(P) = \sum_{\gcd(m,n)=1} \frac{1}{n} \sum_{0 \leq r \leq n-1} \chi\left((mT - P)r\right) = \begin{cases} 1 & \text{if } \operatorname{ord}_E(P) = n, \\ 0 & \text{if } \operatorname{ord}_E(P) \neq n, \end{cases} \tag{10}$$

where $n = \#E(\mathbb{F}_p)$ is the order of the rational group of points.

Proof. As the index $m \geq 1$ ranges over the integers relatively prime to n, the element $mT \in E(\mathbb{F}_p)$ ranges over the elliptic primitive points. Ergo, the linear equation

$$mT - P = \mathcal{O}, \tag{11}$$

where $P, T \in E(\mathbb{F}_p)$ are fixed points, $\mathcal{O}$ is the identity point, and m is a variable such that $0 \leq m \leq n-1$, and $\gcd(m, n) = 1$, has a solution if and only if the fixed point $P \in E(\mathbb{F}_p)$ is an elliptic primitive point. Next, replace $\chi(t) = e^{i2\pi t/n}$ to obtain

$$\Psi_E(P) = \sum_{\gcd(m,n)=1} \frac{1}{n} \sum_{0 \leq r \leq n-1} e^{i2\pi \log_T(mT-P)r/n} = \begin{cases} 1 & \text{if } \operatorname{ord}_E(P) = n, \\ 0 & \text{if } \operatorname{ord}_E(P) \neq n, \end{cases} \tag{12}$$

This follows from the geometric series identity $\sum_{0 \leq k \leq N-1} w^k = (w^N - 1)/(w - 1)$ with $w \neq 1$, applied to the inner sum. ■

3 Primes In Short Intervals

There are many unconditional results for the existence of primes in short intervals $[x, x + y]$ of subsquareroot length $y \leq x^{1/2}$ for almost all large numbers $x \geq 1$. One of the earliest appears to be the Selberg result for $y = x^{19/77}$ with $x \leq X$ and $O\left(X(\log X)^2\right)$ exceptions, see [24, Theorem 4]. Recent improvements based on different analytic methods are given in [32], [10], et alii. One of these results, but not the best, has the following claim. This result involves the weighted prime indicator function (vonMangoldt), which is defined by

$$\Lambda(n) = \begin{cases} \log n & \text{if } n = p^k, k \geq 1, \\ 0 & \text{if } n \neq p^k, k \geq 1, \end{cases} \tag{13}$$

where $p^k, k \geq 1$ is a prime power.

Theorem 3.1. ([10]) *Given $\varepsilon > 0$, almost all intervals of the form $[x - x^{1/6+\varepsilon}, x]$ contains primes except for a set of $x \in [X, 2X]$ with measure $O\left(X(\log X)^{C-1}\right)$, and $C > 1$ constant. Moreover,*

$$\sum_{x-y \leq n \leq x} \Lambda(n) > \frac{y}{2}. \tag{14}$$

Proof. Based on zero density theorems, *ibidem*, p. 190. ■

An introduction to zero density theorems and its application to primes in short intervals is given in [16, p. 98], [12, p. 264], et cetera.

The same result works for larger intervals $[x - x^{\theta+\varepsilon}, x]$, where $\theta \in (1/6, 1/2)$ and $(1-\theta)C < 2$, with exceptions $O\left(X(\log X)^{C-1}\right), C > 1$.

4 Evaluation Of The Main Term

A lower bound for the main term $M(x)$ in the proof of Theorem 1.1 is evaluated here.

Lemma 4.1. *Let $x \geq 1$ be a large number, and let $p \in [x, 2x]$ be prime. Then,*

$$\sum_{x\leq p\leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p}\leq n\leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \sum_{\gcd(m,n)=1} \frac{1}{n} \gg \frac{x}{\log^3 x}\left(1 + O\left(\frac{x}{\log x}\right)\right). \tag{15}$$

Proof. The Euler phi function $\varphi(n) = \#\{1 \leq m < n : \gcd(m, n) = 1\}$, has the lower bound $\varphi(n)/n \gg 1/\log n$ for any integer $n \geq 1$, better estimates are proved in [6, p. 118], [29, p. 55], et cetera. Thus, main term can be rewritten in the form

$$\begin{aligned} M(x) &= \sum_{x\leq p\leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p}\leq n\leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \sum_{\gcd(m,n)=1} \frac{1}{n} \\ &= \sum_{x\leq p\leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p}\leq n\leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n}\frac{\varphi(n)}{n} \qquad (16) \\ &\gg \sum_{x\leq p\leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p}\leq n\leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log^2 n} \\ &\gg \sum_{x\leq p\leq 2x} \frac{1}{4\sqrt{p}} \cdot \frac{1}{\log^2 p} \sum_{p-2\sqrt{p}\leq n\leq p+2\sqrt{p}} \Lambda(n). \qquad (17) \end{aligned}$$

Now, observe that as the prime $p \in [x, 2x]$ varies, the number of short intervals $[p - 2\sqrt{p}, p + 2\sqrt{p}]$ is the same as the number of primes in the interval $[x, 2x]$, namely,

$$\pi(2x) - \pi(x) = \frac{x}{\log x}\left(1 + O\left(\frac{1}{\log x}\right)\right) > \frac{x}{2\log x} \tag{18}$$

for large $x \geq 1$. By Theorem 3.1, the finite sum over the short interval satisfies

$$\sum_{p-2\sqrt{p}\leq n\leq p+2\sqrt{p}} \Lambda(n) > \frac{4\sqrt{p}}{2}, \tag{19}$$

with $O\left(x\log^{-C} x\right)$ exceptions $p \in [x, 2x]$, where $1 < C < 4$. Take $C = 3 + \varepsilon$, where $\varepsilon > 0$ is a small number. Then, the number of exceptions is small in comparison to the number of intervals $\pi(2x) - \pi(x) > x/2\log x$ for large $x \geq 1$. Hence, an

application of this Theorem yields

$$\begin{aligned}\sum_{x\le p\le 2x}\frac{1}{4\sqrt{p}}\cdot\frac{1}{\log^2 p}\sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}}\Lambda(n) &\gg \sum_{x\le p\le 2x}\frac{1}{4\sqrt{p}}\cdot\frac{1}{\log^2 p}\cdot\left(\frac{4\sqrt{p}}{2}\right)+O\left(\frac{x}{\log^{3+\varepsilon} x}\right)\\ &\gg \frac{1}{\log^2 x}\sum_{x\le p\le 2x}1+O\left(\frac{x}{\log^{3+\varepsilon} x}\right) \qquad (20)\\ &\gg \frac{1}{\log^2 x}\cdot\frac{x}{\log x}\left(1+O\left(\frac{1}{\log^{\varepsilon} x}\right)\right),\end{aligned}$$

where all the errors terms are absorbed into one term. ■

Remark 4.1. The exceptional intervals $[p-2\sqrt{p}, p+2\sqrt{p}]$, where $p\in[x,2x]$, contain fewer primes, that is,

$$\sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}}\Lambda(n)=o(\sqrt{p}). \qquad (21)$$

This shortfalls is accounted for in the correction term

$$\begin{aligned}\sum_{x\le p\le 2x}\frac{1}{4\sqrt{p}}\cdot\frac{1}{\log^2 p}\sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}}\Lambda(n) &= o\left(\sum_{x\le p\le 2x}\frac{1}{4\sqrt{p}}\cdot\frac{1}{\log^2 p}\cdot\sqrt{p}\right)\\ &= O\left(\frac{x}{\log^C x}\right) \qquad (22)\end{aligned}$$

in the previous calculation.

5 Estimate For The Error Term

The analysis of an upper bound for the error term $E(x)$, which occurs in the proof of Theorem 1.1, is split into two parts. The first part in Lemma 5.1 is an estimate for the triple inner sum. And the final upper bound is assembled in Lemma 5.2.

Lemma 5.1. *Let E be a nonsingular elliptic curve over rational number, let $P\in E(\mathbb{Q})$ be a point of infinite order. Let $x\ge 1$ be a large number. For each prime $p\ge 3$, fix a primitive point T, and suppose that $P\in E(\mathbb{F}_p)$ is not a primitive point for all primes $p\ge 2$, then*

$$\sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}}\frac{\Lambda(n)}{n\log n}\sum_{\gcd(m,n)=1}\sum_{1\le r<n}\chi((mT-P)r)\le 2. \qquad (23)$$

Proof. Let $\log_T : E(\mathbb{F}_p)\to\mathbb{Z}_n$ be the discrete logarithm function with respect to the fixed primitive point T, defined by $\log_T(mT)=m$, $\log_T(P)=k$, and $\log_T(\mathcal{O})=0$. Then, the nontrivial additive character evaluates to

$$\chi(rmT)=e^{\frac{i2\pi}{n}\log_T(rmT)}=e^{i2\pi rm/n}, \qquad (24)$$

and

$$\chi(-rP) = e^{\frac{i2\pi}{n}\log_T(-rP)} = e^{-i2\pi rk/n}, \tag{25}$$

respectively. To derive a sharp upper bound, rearrange the inner double sum as a product

$$\begin{aligned} T(p) &= \sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}} \frac{\Lambda(n)}{n\log n} \sum_{\gcd(m,n)=1,\ 1\le r<n} \chi((mT-P)r) \\ &= \sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}} \frac{\Lambda(n)}{n\log n} \sum_{1\le r<n} \chi(-rP) \sum_{\gcd(m,n)=1} \chi(rmT) \\ &= \sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}} \frac{\Lambda(n)}{n\log n} \left(\sum_{1\le r<n} e^{-i2\pi rk/n}\right)\left(\sum_{\gcd(m,n)=1} e^{i2\pi rm/n}\right). \end{aligned} \tag{26}$$

The hypothesis $mT - P \neq \mathcal{O}$ for $m \geq 1$ such that $\gcd(m, n) = 1$ implies that the two inner sums in (26) are complete geometric series, except for the terms for $m = 0$ and $r = 0$. Moreover, since $n \geq 2$ is a prime, the two geometric sums have the exact evaluations

$$U_n = \sum_{0<r\le n-1} e^{-i2\pi rk/n} = -1 \qquad \text{and} \qquad V_n = \sum_{\gcd(m,n)=1} e^{i2\pi rm/n} = -1 \tag{27}$$

for $1 \leq k < n$, and $1 \leq r < n$ respectively.

Remark 5.1. The evaluation of the finite sum $U_n = U_n(k) = -1$ is independnet of $k \geq 1$ because $n \geq 2$ is prime and $1 \leq k < n$. Similarly, the evaluation of the finite sum $V_n = V_n(r) = -1$ is independnet of $r \geq 1$ because $n \geq 2$ is prime and $1 \leq r < n$.

Therefore, it reduces to

$$\begin{aligned} T(p) &= \sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}} \frac{\Lambda(n)}{n\log n} \\ &\le \frac{1}{\log p} \sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}} \frac{\Lambda(n)}{n} \\ &\le \frac{1}{\log p} \sum_{n\le p+2\sqrt{p}} \frac{\Lambda(n)}{n} \\ &\le 2. \end{aligned} \tag{28}$$

The last line uses the finite sum $\sum_{n\le x} \Lambda(n)/n \ll \log x$, refer to [23, Theorem 2.7] for additional information. ■

Lemma 5.2. *Let E be a nonsingular elliptic curve over rational number, let $P \in E(\mathbb{Q})$ be a point of infinite order. For each large prime $p \geq 3$, fix a primitive point T, and suppose that $P \in E(\mathbb{F}_p)$ is not a primitive point for all primes $p \geq 2$, then*

$$\sum_{x\le p\le 2x} \sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}} \frac{\Lambda(n)}{n\log n} \sum_{\gcd(m,n)=1,\ 1\le r<n} \chi((mT-P)r) = O(x^{1/2}). \tag{29}$$

Proof. By assumption $P \in E(\mathbb{F}_p)$ is not a primitive point. Hence, the linear equation $mT - P = \mathcal{O}$ has no solution $m \in \{m : \gcd(m,n) = 1\}$. This implies that the discrete logarithm $\log_T(mT - P) \neq 0$, and $\sum_{1 \leq r < n} \chi((mT-P)r) = -1$. This in turns yields

$$\begin{aligned} |E(x)| &= \left| \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{n \log n} \sum_{\gcd(m,n)=1,\ 1 \leq r < n} \chi((mT-P)r) \right| \\ &\leq \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{n \log n} \sum_{\gcd(m,n)=1} 1 \qquad (30) \\ &\ll \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{2 \log n} \\ &\ll \frac{x}{\log^3 x} + O\left(\frac{x}{\log^4 x}\right). \end{aligned}$$

Here, the inequality $(1/n) \sum_{\gcd(m,n)=1} 1 = \varphi(n)/n \leq 1/2$ for all integers $n \geq 1$ was used in the second line. Hence, there is a nontrivial upper bound for the error term. To derive a sharper upper bound, take absolute value, and apply Lemma 5.1 to the inner triple sum to obtain this:

$$\begin{aligned} |E(x)| &\leq \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \left| \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{n \log n} \sum_{\gcd(m,n)=1,\ 1 \leq r < n} \chi((mT-P)r) \right| \\ &\leq \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} (2) \\ &\leq \frac{1}{\sqrt{x}} \sum_{x \leq p \leq 2x} 1 \qquad (31) \\ &= O\left(x^{1/2}\right), \end{aligned}$$

The last line uses the trivial estimate $\sum_{x \leq p \leq 2x} 1 \leq x$. ■

6 Elliptic Divisors

The divisor $\operatorname{div}(f) = \gcd(f(\mathbb{Z}))$ of a polynomial $f(x)$ is the greatest common divisor of all its values over the integers, confer [7, p. 395]. Basically, the same concept extents to the setting of elliptic groups of prime orders, but it is significantly more complex.

Definition 6.1. Let $\mathcal{O}_\mathcal{K}$ be the ring of integers of a quadratic numbers field $\mathcal{K}$. The elliptic divisor is an integer $d_E \geq 1$ defined by

$$d_E = \gcd\left(\{\#E(\mathbb{F}_p) : p \geq 2 \text{ and } p \text{ splits in } \mathcal{O}_\mathcal{K}\}\right). \qquad (32)$$

Considerable works, [3], [22], [11], [14], have gone into determining the elliptic divisors for certain classes of elliptic curves.

Theorem 6.1. ([14, Proposition 1]) *The divisor of an elliptic curve* $E : y^2 = x^3 + ax + b$ *over the rational numbers* $\mathbb{Q}$ *with complex multiplication by* $\mathbb{Q}(\sqrt{D})$ *and conductor* N *satisfies* $d_E|24$. *The complete list, with* $c, m \in \mathbb{Z} - \{0\}$, *is the following.*

D	(a,b)	d_E	D	(a,b)	d_E
-3	$(0,m)$	1	-7	$(-140c^2, -784c^3)$	4
-3	$(0,m^2);(0,-27m^2)$	3	-8	$(-30c^2, -56c^3)$	2
-3	$(0,m^3)$	4	-11	$(-1056c^2, -13552c^3)$	1
-3	$(0,c^6);(0,27c^6)$	12	-19	$(-608c^2, -5776c^3)$	1
-4	$(m,0)$	2	-43	$(-13760c^2, -621264c^3)$	1
-4	$(m^2,0);(-m^2,0)$	4	-67	$(-117920c^2, -15585808c^3)$	1
-4	$(-c^4,0);(4c^4,0)$	8	-163	$(-34790720c^2, -78984748304c^3)$	1

7 Densities Expressions

The product expression appearing in Conjecture 1.1, id est,

$$P_0 = \prod_{p \geq 2} \left(1 - \frac{p^2 - p - 1}{(p-1)^3(p+1)}\right) \approx 0.505166168239435774, \tag{33}$$

is the basic the average density of prime orders, it was proved in [17], and very recently other proofs are given in [1, Theorem 1], [15], [21], et alii. The actual density has a slight dependence on the elliptic curve E and the point P. The determination of the dependence is classified into several cases depending on the torsion groups $E(\mathbb{Q})_{\text{tors}}$, and other parameters.

Lemma 7.1. ([30, Proposition 4.2]). *Let* $E : f(x,y) = 0$ *be a Serre curve over the rational numbers. Let* D *be the discriminant of the numbers field* $\mathbb{Q}(\sqrt{\Delta})$, *where* Δ *is the discriminant of any Weierstrass model of* E *over* $\mathbb{Q}$. *If* $d_E = 1$, *then*

$$\delta(1,E) = \begin{cases} \left(1 + \prod_{q|D} \frac{1}{q^3 - 2q^2 - q + 3}\right) \prod_{p \geq 2} \left(1 - \frac{p^2 - p - 1}{(p-1)^3(p+1)}\right) & \text{if } D \equiv 1 \bmod 4; \\ \prod_{p \geq 2} \left(1 - \frac{p^2 - p - 1}{(p-1)^3(p+1)}\right) & \text{if } D \equiv 0 \bmod 4. \end{cases} \tag{34}$$

8 Elliptic Brun Constant

The upper bound

$$\begin{aligned} \pi(x,E) &= \#\{p \leq x : p \not| \Delta \text{ and } \#E(\mathbb{F}_p) = \text{prime}\} \\ &\ll \frac{x}{(\log x)(\log\log\log x)} \end{aligned} \tag{35}$$

for elliptic curves with complex multiplication was proved in [3, Proposition 7]. The same upper bound for elliptic curves without complex multiplication was proved in [31, Theorem 1.3]. An improved version for all elliptic curves follows easily from methods used here.

Lemma 8.1. *For any large number $x \geq 1$ and any elliptic curves $E : f(X, Y) = 0$ of discriminant $\Delta \neq 0$,*

$$\pi(x, E) \leq \frac{6x}{\log^2 x}. \tag{36}$$

Proof. The number of such elliptic primitive primes has the asymptotic formula

$$\begin{aligned} \pi(x, E) &= \sum_{\substack{p \leq x \\ \mathrm{ord}_E(P)=n \text{ prime}}} 1 \qquad (37) \\ &= \sum_{p \leq x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \cdot \Psi_E(P), \end{aligned}$$

where $\Psi_E(P)$ is the charateritic function of primitive points $P \in E(\mathbb{Q})$. This is obtained from the summation of the elliptic primitive primes density function over the interval $[1, x]$, see (48).

Since $\Psi_E(P) = 0, 1$, the previous equation has the upper bound

$$\begin{aligned} \pi(x, E) &\leq \sum_{p \leq x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \\ &\leq \sum_{p \leq x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq q \leq p+2\sqrt{p}} 1, \end{aligned}$$

where q ranges over the primes in the short interval $[p - 2\sqrt{p}, p + 2\sqrt{p}]$. The inner sum is estimated using either the explicit formula or Brun-Titchmarsh theorem. The later result states that the number of primes p in the short interval $[x, x + 4\sqrt{x}]$ satisfies the inequality

$$\pi(x + 4\sqrt{x}) - \pi(x) \leq \frac{3 \cdot 4\sqrt{x}}{\log x}, \tag{38}$$

see [12, p. 167], [23, Theorem 3.9], and [29, p. 83], and similar references. Replacing (38) into (38) yields

$$\begin{aligned} \sum_{p \leq x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq q \leq p+2\sqrt{p}} 1 &\leq \sum_{p \leq x} \frac{1}{4\sqrt{p}} \left(\frac{3 \cdot 4\sqrt{p}}{\log p} \right) \qquad (39) \\ &\leq 3 \sum_{p \leq x} \frac{1}{\log p} \\ &\leq 6 \frac{x}{\log^2 x}. \end{aligned}$$

p	3	13	19	61	67	73	139	163	211	331
n	3	19	13	61	73	81	163	139	199	331
p	349	541	547	571	613	661	757	829	877	
n	313	571	571	541	661	613	787	823	937	

Table 1: Prime Orders $n = \#E(\mathbb{F}_p)$ modulo p for $y^2 = x^3 + 2$.

The last inequality follows by partial summation and the prime number theorem $\pi(x) = x/\log x + O(x/\log^2 x)$. ■

This result facilitates the calculations of a new collection of constants associated with elliptic curves.

Corollary 8.1. *For any elliptic curve E, the elliptic Brun constant*

$$\sum_{\substack{p\geq 2\\ \#E(\mathbb{F}_p)=prime}} \frac{1}{p} < \infty \tag{40}$$

converges.

Proof. Use the prime counting measure $\pi(x, E) \leq 6x/\log^2 x + O(x/\log^3 x)$ in Lemma 8.1 to evaluate the infinite sum

$$\begin{aligned} \sum_{\substack{p\geq 2\\ \#E(\mathbb{F}_p)=\text{prime}}} \frac{1}{p} &= \int_2^\infty \frac{1}{t} d\pi(t, E) \\ &= O(1) + \int_2^\infty \frac{\pi(t, E)}{t^2} dt \\ &< \infty \end{aligned} \tag{41}$$

as claimed. ■

The elliptic Brun constants and the numerical data for the prime orders of a few elliptic curves were compiled. The last example shows the highest density of prime orders. Accorddingly, it has the largest constant.

Example 8.1. The nonsingular Bachet elliptic curve $E : y^2 = x^3 + 2$ over the rational numbers has complex multiplication by $\mathbb{Z}[\rho]$, and nonzero rank $\text{rk}(E) = 1$. The data for $p \leq 1000$ with prime orders n are listed on the Table 8.1; and the elliptic Brun constant is

$$\sum_{\substack{p\geq 2\\ \#E(\mathbb{F}_p)=\text{prime}}} \frac{1}{p} = .520067922\ldots. \tag{42}$$

p	3	7	97	103	181	271	313	367	409	487
n	3	7	97	107	163	293	331	383	397	499
p	883	967								
n	853	941								

Table 2: Prime Orders $n = \#E(\mathbb{F}_p)$ modulo p for $y^2 = x^3 + 6x - 2$.

p	5	7	11	19	43	67	163	211	283	331
n	2	2	3	5	11	17	41	53	71	83
p	523	547	691	787	907					
n	131	137	173	197	227					

Table 3: Prime Orders $n/4 = \#E(\mathbb{F}_p)/4$ modulo p for $y^2 = x^3 - x$.

Example 8.2. The nonsingular elliptic curve $E : y^2 = x^3 + 6x - 2$ over the rational numbers has no complex multiplication and zero rank $\mathrm{rk}(E) = 0$. The data for $p \leq 1000$ with prime orders n are listed on the Table 8.2; and the elliptic Brun constant is

$$\sum_{\substack{p\geq 2 \\ \#E(\mathbb{F}_p)=\text{prime}}} \frac{1}{p} = .186641187\ldots. \tag{43}$$

Example 8.3. The nonsingular elliptic curve $E : y^2 = x^3 - x$ over the rational numbers has complex multiplication by $\mathbb{Z}[\rho]$, and nonzero rank $\mathrm{rk}(E) = 0$. The data for $p \leq 1000$ with prime orders $n/4$ are listed on the Table 8.3; and the elliptic Brun constant is

$$\sum_{\substack{p\geq 2 \\ \#E(\mathbb{F}_p)/4=\text{prime}}} \frac{1}{p} = .549568584\ldots. \tag{44}$$

Example 8.4. The nonsingular elliptic curve $E : y^2 = x^3 - x$ over the rational numbers has complex multiplication by $\mathbb{Z}[\rho]$, and nonzero rank $\mathrm{rk}(E) = 0$. The data for $p \leq 1000$ with prime orders $n/8$ are listed on the Table 8.4; and the elliptic Brun constant is

$$\sum_{\substack{p\geq 2 \\ \#E(\mathbb{F}_p)/8=\text{prime}}} \frac{1}{p} = .2067391731\ldots. \tag{45}$$

Example 8.5. The nonsingular Bachet elliptic curve $E : y^2 = x^3 + 1$ over the rational numbers has complex multiplication by $\mathbb{Z}[\rho]$, and nonzero rank $\mathrm{rk}(E) =?1$. The data for $p \leq 1000$ with prime orders $n/12$ are listed on the Table 8.5; and the elliptic Brun constant is

$$\sum_{\substack{p\geq 2 \\ \#E(\mathbb{F}_p)/12=\text{prime}}} \frac{1}{p} = .5495685884\ldots. \tag{46}$$

p	17	23	29	37	53	101	103	109	149	151
n	2	3	5	5	5	13	13	13	17	19
p	157	277	293	317	389	487	541	631	661	701
n	17	37	37	41	37	53	61	73	79	89
p	757	773	797	821	823	829	853			
n	97	101	97	109	103	97	101			

Table 4: Prime Orders $n/8 = \#E(\mathbb{F}_p)/8$ modulo p for $y^2 = x^3 - x$.

p	31	43	59	67	73	79	97	103	131	139
n	3	3	5	7	7	7	7	7	11	13
p	151	163	181	199	227	241	337	367	379	409
n	13	13	13	19	19	19	31	31	31	31
p	421	443	463	487	491	523	563	709	751	787
n	37	37	37	37	41	43	47	61	67	61
p	823	829	859	883	907	947	967	991		
n	73	73	67	73	79	79	79	79		

Table 5: Prime Orders $n/12 = \#E(\mathbb{F}_p)/12$ modulo p for $y^2 = x^3 + 1$.

9 Prime Orders n

The characteristic function for primitive points in the group of points $E(\mathbb{F}_p)$ of an elliptic curve $E : f(X, Y) = 0$ has the representation

$$\Psi_E(P) = \begin{cases} 1 & \text{if } \operatorname{ord}_E(P) = n, \\ 0 & \text{if } \operatorname{ord}_E(P) \neq n. \end{cases} \tag{47}$$

The parameter $n = \#E(\mathbb{F}_p)$ is the size of the group of points, and the exact formula for $\Psi_E(P)$ is given in Lemma 2.5.

Since each order n is unique, the weighted sum

$$\frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \cdot \Psi_E(P) = \begin{cases} \dfrac{1}{4\sqrt{p}} \cdot \dfrac{\Lambda(n)}{\log n} & \text{if } \operatorname{ord}_E(P) = n \text{ and } n = q^k, \\ 0 & \text{if } \operatorname{ord}_E(P) \neq n \text{ or } n \neq q^k, \end{cases} \tag{48}$$

where $n = q^k, k \geq 1$, is a prime power, is a discrete measure for the density of elliptic primitive primes $p \geq 2$ such that $P \in E(\overline{\mathbb{Q}})$ is a primitive point of prime power order $n \in [p - 2\sqrt{p}, p + 2\sqrt{p}]$.

Proof. (Theorem 1.1). Let $\langle P \rangle = E(\mathbb{F}_p)$ for at least one large prime $p \leq x_0$, and let $x \geq x_0 \geq 1$ be a large number. Suppose that $P \notin E(\mathbb{Q})_{\text{tors}}$ is not a primitive point

of prime order $\mathrm{ord}_E(P) = n$ in $E(\mathbb{F}_p)$ for all primes $p \geq x$. Then, the sum of the elliptic primes measure over the short interval $[x, 2x]$ vanishes. Id est,

$$0 = \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \cdot \Psi_E(P). \tag{49}$$

Replacing the characteristic function, Lemma 2.5, and expanding the nonexistence equation (49) yield

$$\begin{aligned}
0 &= \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \cdot \Psi_E(P) \\
&= \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \left(\sum_{\gcd(m,n)=1} \frac{1}{n} \sum_{0 \leq r \leq n-1} \chi((mT-P)r) \right) \\
&= \delta(d_E, E) \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \sum_{\gcd(m,n)=1} \frac{1}{n} \\
&\qquad + \sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \sum_{\gcd(m,n)=1} \frac{1}{n} \sum_{1 \leq r \leq n-1} \chi((mT-P)r) \\
&= \delta(d_E, E) M(x) + E(x),
\end{aligned} \tag{50}$$

where $\delta(d_E, E) \geq 0$ is a constant depending on both the fixed elliptic curve $E : f(X, Y) = 0$ and the integer divisor d_E.

The main term $M(x)$ is determined by a finite sum over the principal character $\chi = 1$, and the error term $E(x)$ is determined by a finite sum over the nontrivial multiplicative characters $\chi \neq 1$.

Applying Lemma 4.1 to the main term, and Lemma 5.2 to the error term yield

$$\begin{aligned}
\sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \cdot \Psi_E(P) &= \delta(d_E, E) M(x) + E(x) \\
&\gg \delta(d_E, E) \frac{x}{\log^3 x} \left(1 + O\left(\frac{x}{\log x} \right) \right) \\
&\qquad + O\left(x^{1/2} \right) \\
&\gg \delta(d_E, E) \frac{x}{\log^3 x} \left(1 + O\left(\frac{x}{\log x} \right) \right).
\end{aligned} \tag{51}$$

But, if $\delta(d_E, E) > 0$, the expression

$$\begin{aligned}
\sum_{x \leq p \leq 2x} \frac{1}{4\sqrt{p}} \sum_{p-2\sqrt{p} \leq n \leq p+2\sqrt{p}} \frac{\Lambda(n)}{\log n} \cdot \Psi_E(P) &\gg \delta(d_E, E) \frac{x}{\log^3 x} \left(1 + O\left(\frac{x}{\log x} \right) \right) \\
&> 0,
\end{aligned} \tag{52}$$

contradicts the hypothesis (49) for all large numbers $x \geq x_0$. Ergo, there are infinitely many primes $p \geq x$ such that a fixed elliptic curve of rank $\mathrm{rk}(E) > 0$ with a

primitive point P of infinite order, for which the corresponding groups $E(\mathbb{F}_p)$ have prime orders. Lastly, the number of such elliptic primitive primes has the asymptotic formula

$$\begin{aligned}\pi(x,E) &= \sum_{\substack{p\le x\\ \mathrm{ord}_E(P)=n \text{ prime}}} 1 \\ &= \sum_{p\le x}\frac{1}{4\sqrt{p}}\sum_{p-2\sqrt{p}\le n\le p+2\sqrt{p}}\frac{\Lambda(n)}{\log n}\cdot\Psi_E(P) \qquad (53)\\ &\ge \delta(d_E,E)\frac{x}{\log^3 x}\left(1+O\left(\frac{x}{\log x}\right)\right),\end{aligned}$$

which is obtained from the summation of the elliptic primitive primes density function over the interval $[1,x]$. ■

The exact evaluation

$$\pi(x,E)=\delta(d_E,E)\frac{x}{\log^2 x}\left(1+O\left(\frac{x}{\log x}\right)\right), \qquad (54)$$

as claimed in Conjecture 1.1, might requires more precise information for primes in short intervals $[p-2\sqrt{p},p-2\sqrt{p}]$.

10 Examples Of Elliptic Curves

The densities of several elliptic curves have been computed by several authors. Extensive calculations for some specific densities are given in [30].

Example 10.1. The nonsingular Bachet elliptic curve $E: y^2=x^3+2$ over the rational numbers has complex multiplication by $\mathbb{Z}[\rho]$, and nonzero rank $\mathrm{rk}(E)=1$. It is listed as 1728.n4 in [20]. The numerical data shows that $\#E(\mathbb{F}_p)=n$ is prime for at least one prime, see Table 8.1. Hence, by Theorem 1.1, the corresponding group of $\mathbb{F}_p$-rational points $\#E(\mathbb{F}_p)$ has prime orders $n=\#E(\mathbb{F}_p)$ for infinitely many primes $p\ge 3$.

Since $\Delta=-2^6\cdot 3^3$, the discriminant of the quadratic field $\mathbb{Q}(\sqrt{\Delta})$ is $D=-3$. Moreover, the integer divisor $d_E=1$ since $\#E(\mathbb{F}_p)$ is prime for at least one prime. Thus, applying Lemma 7.1, gives the natural density

$$\delta(1,E)=\frac{10}{9}P_0\approx 0.561295742488261971297938 5\ldots, \qquad (55)$$

The predicted number of elliptic primes $p\nmid 6N$ such that $\#E(\mathbb{F}_p)$ is prime has the asymptotic formula

$$\pi(x,E)=\delta(1,E)\int_2^x\frac{1}{\log(t+1)}\frac{dt}{t}. \qquad (56)$$

Invariant	**Value**		
Discriminant	Δ	$=$	$-16(4a^3+27b^2) = -1728$
Conductor	N	$=$	1728
j-Invariant	$j(E)$	$=$	$(-48b)^3/\Delta = 0$
Rank	$\mathrm{rk}(E)$	$=$	1
Special L-Value	$L^{'}(E,1)$	$\approx$	2.82785747365
Regulator	R	$=$	.754576
Real Period	Ω	$=$	5.24411510858
Torsion Group	$E(\mathbb{Q})_{\mathrm{tors}}$	$\cong$	$\{\mathcal{O}\}$
Integral Points	$E(\mathbb{Z})$	$=$	$\{\mathcal{O},(-1,1);(-1,1)\}$
Rational Group	$E(\mathbb{Q})$	$=$	$\mathbb{Z}$
Endomorphims Group	$End(E)$	$=$	$\mathbb{Z}[(1+\sqrt{-3})/2]$, CM
Integer Divisor	d_E	$=$	1

Table 6: Data for $y^2 = x^3 + 2$.

A lower bound for the counting function is

$$\pi(x,E) \geq \delta(1,E)\frac{x}{\log^3 x}\left(1+O\left(\frac{1}{\log x}\right)\right), \tag{57}$$

see Theorem 1.1.

The associated weight $k=2$ cusp form, and L-function are

$$f(s) = \sum_{n\geq 1} a_n q^n = q - q^7 - 5q^{13} + 7q^{19} + \cdots, \tag{58}$$

and

$$\begin{aligned} L(s) &= \sum_{n\geq 1}\frac{a_n}{n^s} \\ &= \prod_{p|N}\left(1-\frac{a_p}{p^s}\right)^{-1}\prod_{p\nmid N}\left(1-\frac{a_p}{p^s}+\frac{1}{p^{2s-1}}\right)^{-1} \\ &= 1-\frac{1}{7^s}-\frac{5}{13^s}+\frac{7}{19^s}+\cdots, \end{aligned} \tag{59}$$

where $q = e^{i2\pi}$, respectively. The coefficients are generated using $a_p = p+1-\#E(\mathbb{F}_p)$, and the formulas

1. $a_{pq} = a_p a_q$ if $\gcd(p,q)=1$;
2. $a_{p^{n+1}} = a_{p^n}a_p - p a_{p^{n-1}}$ if $n \geq 2$.

The corresponding functional equation is

$$\Lambda(s) = \left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma(s)L(s) \qquad \text{and} \qquad \Lambda(s) = \Lambda(2-s), \tag{60}$$

where $N = 1728$, see [18, p. 80].

Invariant	**Value**		
Discriminant	Δ	$=$	$-16(4a^3+27b^2) = -2^6 \cdot 3^5$
Conductor	N	$=$	$2^6 \cdot 3^3$
j-Invariant	$j(E)$	$=$	$(-48b)^3/\Delta = 2^9 \cdot 3$
Rank	$\mathrm{rk}(E)$	$=$	0
Special L-Value	$L(E,1)$	$\approx$	2.24402797314
Regulator	R	$=$	1
Real Period	Ω	$=$	2.2440797314
Torsion Group	$E(\mathbb{Q})_{\mathrm{tors}}$	$=$	$\{\mathcal{O}\}$
Integral Points	$E(\mathbb{Z})$	$=$	$\{\mathcal{O}\}$
Rational Group	$E(\mathbb{Q})$	$=$	$\{\mathcal{O}\}$
Endomorphims Group	$End(E)$	$=$	$\mathbb{Z}$, nonCM
Integer Divisor	d_E	$=$	1

Table 7: Data for $y^2 = x^3 + 6x - 2$.

Example 10.2. The nonsingular elliptic curve $E : y^2 = x^3+6x-2$ over the rational numbers has no complex multiplication and zero rank, it is listed as 1728.w1 in [20]. The numerical data shows that $\#E(\mathbb{F}_p) = n$ is prime for at least one prime, see Table 8.2. Hence, by Theorem 1.1, the corresponding group of $\mathbb{F}_p$-rational points $\#E(\mathbb{F}_p)$ has prime orders $n = \#E(\mathbb{F}_p)$ for infinitely many primes $p \geq 3$.

Since $\Delta = -2^6 \cdot 3^5$, the discriminant of the quadratic field $\mathbb{Q}(\sqrt{\Delta})$ is $D = -3$. Moreover, the integer divisor $d_E = 1$ since $\#E(\mathbb{F}_p)$ is prime for at least one prime. Thus, applying Lemma 7.1, gives the natural density

$$\delta(1,E) = \frac{10}{9}P_0 \approx 0.561295742488261971297938 5\ldots, \tag{61}$$

The predicted number of elliptic primes $p \nmid 6N$ such that $\#E(\mathbb{F}_p)$ is prime has the asymptotic formula

$$\pi(x,E) = \delta(1,E)\int_2^x \frac{1}{\log(t+1)}\frac{dt}{t}. \tag{62}$$

A table for the prime counting function $\pi(1,E)$, for $2 \times 10^7 \leq x \leq 10^9$, and other information on this elliptic curve appears in [30].

A lower bound for the counting function is

$$\pi(x,E) \geq \delta(1,E)\frac{x}{\log^3 x}\left(1 + O\left(\frac{1}{\log x}\right)\right), \tag{63}$$

see Theorem 1.1.

The associated weight $k = 2$ cusp form, and L-function are

$$f(s) = \sum_{n\geq 1} a_n q^n = q + 2q^5 + q^7 + 2q^{11} - q^{13} - 6q^{17} + 5q^{19} + \cdots, \tag{64}$$

and

$$\begin{aligned} L(s) &= \sum_{n\geq 1} \frac{a_n}{n^s} \\ &= \prod_{p|N} \left(1 - \frac{a_p}{p^s}\right)^{-1} \prod_{p\nmid N} \left(1 - \frac{a_p}{p^s} + \frac{1}{p^{2s-1}}\right)^{-1} \\ &= 1 + \frac{2}{5^s} + \frac{1}{7^s} + \frac{2}{11^s} - \frac{1}{13^s} - \frac{6}{17^s} + \frac{5}{19^s} + \cdots, \end{aligned} \tag{65}$$

where $q = e^{i2\pi}$, respectively. The coefficients are generated using $a_p = p + 1 - \#E(\mathbb{F}_p)$, and the formulas

1. $a_{pq} = a_p a_q$ if $\gcd(p, q) = 1$;
2. $a_{p^{n+1}} = a_{p^n} a_p - p a_{p^{n-1}}$ if $n \geq 2$.

The corresponding functional equation is

$$\Lambda(s) = \left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma(s) L(s) \qquad \text{and} \qquad \Lambda(s) = \Lambda(2 - s), \tag{66}$$

where $N = 1728$, see [18, p. 80].

Example 10.3. The nonsingular elliptic curve $E : y^2 = x^3 - x$ over the rational numbers has complex multiplication by $\mathbb{Z}[i]$, and zero rank, it is listed as 32.a3 in [20]. The numerical data shows that $\#E(\mathbb{F}_p)/8 = n$ is prime for at least one prime, see (**??**). Hence, by Theorem 1.1, the corresponding group of $\mathbb{F}_p$-rational points $\#E(\mathbb{F}_p)$ has prime orders $n = \#E(\mathbb{F}_p)$ for infinitely many primes $p \geq 3$.

Since $\Delta = -2^6$, the discriminant of the quadratic field $\mathbb{Q}(\sqrt{\Delta})$ is $D = -4$. Moreover, the integer divisor $d_E = 8$ since $\#E(\mathbb{F}_p)/8$ is prime for at least one prime, see Table 8.4. Thus, Lemma 7.1 is not applicable. The natural density

$$\delta(8, E) = \frac{1}{2} \prod_{p\geq 3} \left(1 - \chi(p) \frac{p^2 - p - 1}{(p - \chi(p))(p - 1)^2}\right) \approx 0.5336675447\ldots, \tag{67}$$

where $\chi(n) = (-1)^{(n-1)/2}$, is computed in [30, Lemma 7.1]. The predicted number of elliptic primes $p \nmid 6N$ such that $\#E(\mathbb{F}_p)$ is prime has the asymptotic formula

$$\pi(x, E) = \delta(8, E) \int_9^x \frac{1}{\log(t + 1) - \log 8} \frac{dt}{\log t}. \tag{68}$$

A table for the prime counting function $\pi(x, E)$, for $2 \times 10^7 \leq x \leq 10^9$, and other information on this elliptic curve appears in [30].

A lower bound for the counting function is

$$\pi(x, E) \geq \delta(8, E) \frac{x}{\log^3 x} \left(1 + O\left(\frac{1}{\log x}\right)\right), \tag{69}$$

Invariant	**Value**		
Discriminant	Δ	$=$	$-16(4a^3+27b^2) = -2^6$
Conductor	N	$=$	2^5
j-Invariant	$j(E)$	$=$	$(-48b)^3/\Delta = 2^6 \cdot 3^3$
Rank	$\mathrm{rk}(E)$	$=$	0
Special L-Value	$L(E,1)$	$\approx$	$.655514388573$
Regulator	R	$=$	1
Real Period	Ω	$=$	5.24411510858
Torsion Group	$E(\mathbb{Q})_{\mathrm{tors}}$	$\cong$	$\mathbb{Z}_2 \times \mathbb{Z}_2$
Integral Points	$E(\mathbb{Z})$	$=$	$\{\mathcal{O}, (\pm 1, 0); (0,0)\}$
Rational Group	$E(\mathbb{Q})$	$=$	$E(\mathbb{Q})_{\mathrm{tors}}$
Endomorphims Group	$End(E)$	$=$	$\mathbb{Z}[i]$, CM
Integer Divisor	d_E	$=$	$2, 4, 8$

Table 8: Data for $y^2 = x^3 - x$.

see Theorem 1.1.

The associated weight $k = 2$ cusp form, and L-function are

$$f(s) = \sum_{n \geq 1} a_n q^n = q + 2q^5 + q^7 + 2q^{11} - q^{13} - 6q^{17} + 5q^{19} + \cdots, \tag{70}$$

and

$$\begin{aligned} L(s) &= \sum_{n \geq 1} \frac{a_n}{n^s} \\ &= \prod_{p|N} \left(1 - \frac{a_p}{p^s}\right)^{-1} \prod_{p \nmid N} \left(1 - \frac{a_p}{p^s} + \frac{1}{p^{2s-1}}\right)^{-1} \\ &= 1 + \frac{2}{5^s} + \frac{1}{7^s} + \frac{2}{11^s} - \frac{1}{13^s} - \frac{6}{17^s} + \frac{5}{19^s} + \cdots, \end{aligned} \tag{71}$$

where $q = e^{i2\pi}$, respectively. The coefficients are generated using $a_p = p + 1 - \#E(\mathbb{F}_p)$, and the formulas

1. $a_{pq} = a_p a_q$ if $\gcd(p, q) = 1$;
2. $a_{p^{n+1}} = a_{p^n} a_p - p a_{p^{n-1}}$ if $n \geq 2$.

The corresponding functional equation is

$$\Lambda(s) = \left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma(s) L(s) \qquad \text{and} \qquad \Lambda(s) = \Lambda(2 - s), \tag{72}$$

where $N = 1728$, see [18, p. 80].

Exercises

Problem 1. Assume random elliptic curve $E : f(x, y) = 0$ has no CM, P is a point of infinite order. Is the integer divisor $d_E < \infty$ bounded? This parameter is bounded for CM elliptic curves, in fact $d_E < 24$, reference: [14, Proposition 1].

Problem 2. Assume the elliptic curve $E : f(x, y) = 0$ has no CM, P is a point of infinite order, and the density $\delta(1, E) > 0$. What is the least prime $p \nmid 6N$ such that the integer divisor $d_E = 1$? Reference: [5, Corollary 1.2].

Problem 3. Fix an elliptic curve $E : y^2 = x^3 + ax + b$ of rank $\mathrm{rk}(E) > 0$, and CM; and P is a point of infinite order. Let the integer divisor $d_E = 4$. Assume the densities $\delta(1, E) > 0$, $\delta(2, E) > 0$, and $\delta(4, E) > 0$ are defined. What is the arithmetic relationship between the densities?

Problem 4. Fix an elliptic curve $E : y^2 = x^3 + ax + b$ of rank $\mathrm{rk}(E) > 0$, and CM; and P is a point of infinite order. Does $\#E(\mathbb{F}_p) = n$ prime for at least one large prime $p \geq 2$ implies that the integer divisor $d_E = 1$?

References

[1] Balog, Antal; Cojocaru, Alina-Carmen; David, Chantal. Average twin prime conjecture for elliptic curves. Amer. J. Math. 133,(2011), no. 5, 1179-1229.

[2] Borosh, I.; Moreno, C. J.; Porta, H. Elliptic curves over finite fields. II. Math. Comput. 29 (1975), 951-964.

[3] Cojocaru, Alina Carmen. Reductions of an elliptic curve with almost prime orders. Acta Arith. 119 (2005), no. 3, 265-289.

[4] Cojocaru, Alina Carmen. Questions about the reductions modulo primes of an elliptic curve. Number theory, 61-79, CRM Proc. Lecture Notes, 36, Amer. Math. Soc., Providence, RI, 2004.

[5] Cojocaru, Alina Carmen. Cyclicity of CM elliptic curves modulo p. Trans. Amer. Math. Soc. 355 (2003), no. 7, 2651-2662.

[6] DeKoning, Analytic Number Theory, AMS. 2012.

[7] Friedlander, John; Iwaniec, Henryk. Opera de cribro. American Mathematical Society Colloquium Publications, 57. American Mathematical Society, Providence, RI, 2010.

[8] Gupta, Rajiv; Murty, M. Ram. Primitive points on elliptic curves. Compositio Math. 58 (1986), no. 1, 13-44.

[9] Gupta, Rajiv; Murty, M. Ram. Cyclicity and generation of points mod p on elliptic curves. Invent. Math. 101 (1990), no. 1, 225-235.

[10] Harman, Glyn. Prime-detecting sieves. London Mathematical Society Monographs Series, 33. Princeton University Press, Princeton, NJ, 2007.

[11] Iwaniec, Henryk; Jimenez Urroz, Jorge. Orders of CM elliptic curves modulo p with at most two primes. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 9 (2010), no. 4, 815-832.

[12] Iwaniec, Henryk; Kowalski, Emmanuel. Analytic number theory. AMS Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004.

[13] Jones, Nathan. Averages of elliptic curve constants. Math. Ann. 345 (2009), no. 3, 685-710.

[14] Jimenez Urroz, Jorge. Almost prime orders of CM elliptic curves modulo p. Algorithmic number theory, 74-87, Lecture Notes in Comput. Sci., 5011, Springer, Berlin, 2008.

[15] Jones, Nathan. Almost all elliptic curves are Serre curves. Trans. Amer. Math. Soc. 362 (2010), no. 3, 1547-1570.

[16] Anatolij A. Karacuba, Basic Analytic Number Theory, Springer-Verlag, 1993.

[17] Koblitz, Neal. Primality of the number of points on an elliptic curve over a finite field. Pacific J. Math. 131 (1988), no. 1, 157-165.

[18] Koblitz, Neal. Introduction to elliptic curves and modular forms. Second edition. Graduate Texts in Mathematics, 97. Springer-Verlag, New York, 1993.

[19] Lang, S.; Trotter, H. Primitive points on elliptic curves. Bull. Amer. Math. Soc. 83 (1977), no. 2, 289-292.

[20] The L-functions and modular forms database, www.lmfdb.org.

[21] Lenstra, H. W., Jr.; Stevenhagen, P.; Moree, P. Character sums for primitive root densities. Math. Proc. Cambridge Philos. Soc. 157 (2014), no. 3, 489-511. And arXiv:1112.4816.

[22] Giulio Meleleo, Questions Related To Primitive Points On Elliptic Curves And Statistics For Biquadratic Curves Over Finite Fields, Thesis, Universita degli Studi, Roma Tre, 2015.

[23] Montgomery, Hugh L.; Vaughan, Robert C. Multiplicative number theory. I. Classical theory. Cambridge University Press, Cambridge, 2007.

[24] Selberg, Atle. On the normal density of primes in small intervals, and the difference between consecutive primes. Arch. Math. Naturvid. 47, (1943). no. 6, 87-105.

[25] Silverman, Joseph H. The arithmetic of elliptic curves. Second edition. Graduate Texts in Mathematics, 106. Springer, Dordrecht, 2009.

[26] Silverman, Joseph H. p-adic properties of division polynomials and elliptic divisibility sequences. Math. Ann. 332 (2005), no. 2, 443-471.

[27] Shparlinski, Igor E.; Voloch, Jose Felipe. Generators of elliptic curves over finite fields. Preprint 2011.

[28] Schmitt, Susanne; Zimmer, Horst G. Elliptic curves. A computational approach. With an appendix by Attila Petho. de Gruyter Studies in Mathematics, 31. Walter de Gruyter & Co., Berlin, 2003.

[29] Tenenbaum, Gerald. Introduction to analytic and probabilistic number theory. Translated from the Third French edition. American Mathematical Society, Rhode Island, 2015.

[30] Zywina, David. A refinement of Koblitz's conjecture, arXiv:0909.5280.

[31] Zywina, David. The Large Sieve and Galois Representations, arXiv:0812.222.

[32] Watt, N. Short intervals almost all containing primes. Acta Arith. 72 (1995), no. 2, 131-167.